



Spectroscopy for Industrial Applications: High-Temperature Processes

Fateev, Alexander; Grosch, Helge; Clausen, Sønnik; Barton, Emma J.; Yurchenko, Sergei N.; Tennyson, Jonathan

Publication date:
2014

Document Version
Peer reviewed version

[Link back to DTU Orbit](#)

Citation (APA):
Fateev, A. (Author), Grosch, H. (Author), Clausen, S. (Author), Barton, E. J. (Author), Yurchenko, S. N. (Author), & Tennyson, J. (Author). (2014). Spectroscopy for Industrial Applications: High-Temperature Processes. Sound/Visual production (digital)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.


Spectroscopy for Industrial Applications: High-Temperature Processes

Alexander Fateev^{*)}, Helge Grosch, Sønnik Clausen (DTU Chemical Engineering, Denmark) and

Emma J Barton, Sergei N Yurchenko, Jonathan Tennyson (Department of Physics and Astronomy, UCL, UK)


^{*)} e-mail: alfa@kt.dtu.dk


DTU Chemical Engineering
Department of Chemical and Biochemical Engineering




Outline

- Hot flow gas cell and FTIR/UV optical set up
- A road to In Situ measurements:
 - NH₃ spectroscopy at high-temperatures: band assignment and spectra modelling
 - NH₃/H₂O field measurements at a pilot scale 6MW gasifier
 - Phenol –major trace gas from PAH's in low temperature gasification
 - Temperature-dependent UV absorption cross-sections
 - Why In Situ measurements are important: comparison with “standard” tools
- How planets meet the Earth
- Conclusions





- 3-zones flow gas cell for corrosive gases;
- No internal windows;
- Stable uniform T-profile ($\pm 1.8^\circ\text{C}$);
- $T_{\text{max}} = 525^\circ\text{C}$
- $L = 33.25\text{ cm}$
- $P = 1\text{ bar}$
- suitable for UV-FIR optical measurements
- [more details: H. Grosch et al. JQSRT 130 \(2013\) 392–399](#)
- FTIR Spectrometer (Agilent 660), 0.09 cm^{-1}
- an IR light source (up to 1500°C)
- UV spectrometer (Acton 250i/CCD), 0.019 nm
- a highly stable D2-lamp








Table 1: Lines assigned to previously observed bands

Band	Lines Assigned This Work				Previously measured			
	J_{\max}	K_{\max}	Frequency Range cm^{-1}	Number of Lines	J_{\max}	K_{\max}	Number of Lines	Reference
ν_4	17	17	1290 - 1868	277	15	15	1663	Cottaz 2000
ν_2	20	20	634 - 1333	385	23	20	177	Yu 2010
$\nu_2+\nu_4-\nu_2$	12	12	1412 - 1818	83	10		384	Cottaz 2001
$2\nu_2$	16	15	1407 - 1870	43	15	15	403	Cottaz 2000
$2\nu_2-\nu_2$	18	18	607 - 1236	180	10	10	32	Singh 1988
$3\nu_2-\nu_2$	12	12	1104 - 1652	18	10		132	Cottaz 2001

500C
1 bar
0.09cm-1

Table 2: Lines assigned to previously unobserved bands* with 10 or more lines assigned in this work.


Band	Lines Assigned This Work			
	J_{\max}	K_{\max}	Frequency Range cm^{-1}	Number of Lines
$\nu_4-\nu_2$	11	11	622 - 1013	20
$2\nu_4-\nu_4$	9	5	1430 - 1792	10
$2\nu_4^2-\nu_4$	8	5	1420 - 1805	10
$\nu_2+\nu_4-\nu_4$	13	13	680 - 1270	77
$3\nu_2-2\nu_2$	14	12	628 - 1455	31
$3\nu_2-3\nu_2$	12	9	628 - 743	12



List of Assigned Lines

BYTe Frequency	Experimental Frequency	Obs - Calc	Upper Quantum Numbers*	Lower Quantum Numbers*
968.761998	968.825639	0.063641	- 0 2 0 0 0 16 16	+ 0 1 0 0 0 16 16
970.332898	970.874628	0.54173	- 0 2 0 0 0 14 3	+ 0 1 0 0 0 13 3
971.871137	971.868991	-0.002146	- 0 1 0 0 0 2 1	+ 0 0 0 0 0 1 1
972.159794	972.456569	0.296775	+ 0 2 0 0 0 14 2	- 0 1 0 0 0 13 2
972.363167	972.60723	0.244063	+ 0 2 0 0 0 17 17	- 0 1 0 0 0 17 17
972.801729	973.330403	0.528674	- 0 2 0 0 0 14 1	+ 0 1 0 0 0 13 1
974.317864	974.354898	0.037034	- 0 1 0 1 0 13 1	+ 0 0 0 1 0 12 1
975.511534	975.530054	0.01852	+ 0 1 0 1 0 13 2	- 0 0 0 1 0 12 2
976.392929	976.449086	0.056157	- 0 2 0 0 0 18 18	+ 0 1 0 0 0 18 18


*Parity $v_1 v_2 v_3 v_4 l_3 l_4 J K$



Can we use BYTe at 500C for practical apps?

- in general a good agreement
- some difficulties with strong line intensities
- some frequency shifts in line positions

More work to do at even higher T (>500C)



From Lab to Field

In Situ measurements on Pyroneer (6MW) gasifier

Pyroneer

Home Sitemap All DONG Energy sites

Applications Gasification process Demonstration plant Research and development About Pyroneer News

Pyroneer

A new Gasolution
- Biomass Gasification

Low Temperature
Gasification



Currently a 6MW demonstration plant is

LT-CFB



Gasification is a method for producing
clean fuels from waste biomass and
therefore has the potential to play a vast

Fuel and ash
partnerships



DONG Energy will supply future

6MW Project Status



Keep up with the status of our 6MW
demonstration plant

- Very complex producer gas composition (CO_2 , H_2O , CO , H_2 , HC , PAH , tars) + particles
- Producer gas is feeded into an industrial burner of a power plant

Why to do it? (examples):

- H_2O (related to mass balance)
- NH_3 (related to NO_x formation)

How?: In Situ IR abs measurements: no gas extraction


- Tough: out of the building on a platform (safety) with limited space (practical issues);
- T_{gas} about 530°C ;
- optical measurements over 30 cm;
- very strong any (UV-IR) light attenuation.

Pyroner 25-06-2013 (30 cm, 530°C , 1.1 bar)

blue: high resolution (0.5cm^{-1})
red: reference H_2O

log10_sb_c1_c0.csv 0.5cm-1
H2O=35% 30cm 530C 1.1 bar

$\text{H}_2\text{O}=35\%/30\text{cm}$





Pyroner 25-06-2013 (30 cm, 530°C , 1.1 bar)

blue: high resolution (0.5cm^{-1})
red: reference NH_3

log10_sb_c1_c0.csv 0.5cm-1
 $\text{NH}_3=0.6\%$ 30cm 530C 1.1 bar

$\text{NH}_3=0.6\%/30\text{cm}$





1. Gas mixing unit

- N_2 (industrial standard)
- molten aromatic crystals in tube
- ⇒ concentration unknown
- admixture of N_2 for different concentration

2. Gas cell and optics

3. Petersen column


- sampling in acetone
- Sampling time 30 min
- analysis with GC/MS

Measurements strategy:


- At each T two phenol concentrations
- At each concentration two sample
- During each sampling three UV spectra and three double concentration determination

- Not too many reference data available even at low T (about 23C)
- An excellent agreement with published data at low T
- Significant changes in the fine structure of the cross-section spectra with T

Low-temperature abs cross-sections: comparison





Abs cross-sections: from 23C to 500C






- Focus on trace gases in low- and high-temperature gasification processes;
- Producer gas issues:
 - corrosion (boilers)
 - reduced gas quality (fuel cells, gas grids)
- Phenol – major trace gas from PAH's in the producer gas (LT-CFB process);
- $T_{\text{gas}} = 300\text{-}500\text{C}$; In Situ UV abs measurements over 3 mm;
- Phenol measurements by various techniques:
 - GC/MS (Petersen column (30 min) 215 ppm ($\pm 5\%$))
 - Gas extraction, 150 C: 407 ppm ($\pm 5\%$) (3 min)
 - In Situ, 306 C (DOAS approach): 7700 ppm ($\pm 10\%$) (3 min)



- Far away planets on a global scale (e.g. exoplanets, stars) and current Earth's problems on a local scale (energy, emissions , taxes)
- Spectroscopy of hot planets and high-temperature processes: the same gases/temperatures of interest;
- DTU's projects about optical measurements in combustion (SO_2 , SO_3 , NH_3 , etc), gasification (trace gases, Cl- compounds) and waste utilization in collaboration with industry (DONG Energy, Vattenfall and Babcock & Wilcox Vølund)
- UCL's and DTU's common PhD/postdocs projects: SO_3/SO_2 and Cl-compounds (KCl, HCl, CH_3Cl , CH_4 , H_2CO)



Conclusions

In general

- You can find a lot inspirations for the work on the Earth
- Different research areas can have the same origin
- Scientists can make industry guys happy

In particular:

- Excellent experimental tools are available for (VUV) UV-FIR optical measurements
- Temperature range can be also negative (e.g. gases at low T)
- New data/lines for NH₃ BYTe extension and development
- New data for phenol
- Try always In Situ and avoid any Ex Situ (extraction) measurements



- To Energinet.dk: projects No. 2013-12027, 2011-1-10622, 2010-1-10422
- To MST.dk
- To DONG Energy and Vattenfall
- To UCL (Prof. Jonathan Tennyson's group)

